

Table S1—Survey results regarding quality-of-life, sleepiness, family impact, and decisional conflict for children without tonsillar hypertrophy in an upper airway clinic comparing those who had and those who had not undergone previous adenotonsillectomy.

mean±SD	Previous T&A (N=42)	No Previous T&A (N=35)	P value
Decisional Conflict Total	11.4±12.0(0-37.5)	9.6±18.2(0-93.75)	0.19
DC Informed	9.5±11.1(0-33.3)	9.8±19.4(0-100)	0.41
DC Values	11.2±13.6(0-41.7)	10.5±18.0(0-83.3)	0.42
DC Support	9.3±14.1 (0-50)	8.1±18.6 (0-100)	
DC Uncertainty	14.1±17.6(0-100)	11.4±20.9(0-100)	0.24
DC Effectiveness	12.5±14.8(0-87.5)	8.8±17.8 (0-87.5)	0.19
PSQ	0.40 ± 0.30	0.30 ± 0.25	0.18
OSA-18	44.5 ± 20.2	40.5 ± 21.6	0.42
Epworth Sleepiness Scale	7.8 ± 5.7	5.9 ± 4.8	0.31
Family Impact Score			
Negative feelings toward parenting	6.9 ± 5.4	8.2 ± 4.2	0.26
Positive feelings toward parenting	15.1 ± 5.1	12.4 ± 5.3	0.11
Negative impact on social life	7.0 ± 6.5	4.8 ± 2.9	0.71
Financial impact	4.5 ± 4.4	4.1 ± 4.7	0.69
Impact on marriage	3.9 ± 3.0	3.9 ± 2.9	0.98
Impact on siblings	4.7 ± 3.6	5.3 ± 3.2	0.68
CollaboRATE score	93.5 ± 11.4	95.4 ± 12.2	0.14

N=number, SD=standard deviation, PedsQL= pediatric quality of life survey, PSQ = pediatric sleep questionnaire; this uses a cutoff of 0.33 to indicate high risk.